



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

character in each species of tree. They are not merely abnormal developments of the bark, but are composed of true woody fibre disposed in annual rings and surrounded by bark, as though they were so many separate centers of growth. Mr. Meehan suggests the following explanation of these excrescences. By experiment it is shown that there is no difference in the first place in any part of the cambium layer or layer of growth, although subsequently the inner part produces wood cells. "The same cell may become permanent tissue or generating tissue, and from the generative tissue may come before the season of growth closes every form of structure known to anatomists, from pure wood to the outermost cuticle of the bark. We know that cell-growth is not always uniform in its operations. The law that changes the outermost series of newly cells into liber need not necessarily operate so exactly as to make them perfect to this end—a few may be thrown off into the liber as generative tissue—and, granting this possibility, we see how the woody granules in the apple bark are formed." In this same way, Mr. Meehan infers, the larger excrescences with rings of growth are formed. In this case "the imperfectly formed liber cells, still retaining their generative power, make a growth the next season, forming a layer of wood and making its own cortical layer, simultaneous with the normal wood growth of the tree, assimilating from the same store of reserve material that the normal growth does."—J. M. C.

WOLFIA.—*Wolfia* in Gray's Manual, page 480 (Ed. of 1869), is described as having "fronds rootless, proliferous from a cleft or funnel-shaped opening at the base, the offspring soon detached." Last summer I watched *Wolfia* pretty closely and with much interest. In plants from my locality the manner of its reproduction seems to be this. I noticed plants that were nearly spheres, others that were oblong, or prolate spheroids, and of these some had indentations about the middle, which varied in depth, until in some of the plants the two portions were merely in contact and these soon went apart, making two perfect plants.—DR. H. C. BEARDSLEE, *Painesville, Ohio*.

Prof. Theo. B. Comstock, of Cornell University, proposes to conduct an "Aquatic Summer School of Natural History." A steamer is to be chartered for four or six weeks to cruise along the "south and west shores of Lakes Erie, Huron and Superior, thence along the north and east shores of the same waters." The route is certainly a tempting one to botanists and the fee, \$125 including everything for 30 days, is so moderate that we have no doubt many will be tempted to make application to join the expedition.

We take pleasure in calling attention to a circular issued by H. Eggert of St. Louis, Mo. It surely presents a rare chance to botanists to obtain sets of the fine and many rare plants found in the neighborhood of St. Louis. We have received a set of Mr. Eggert's plants and have found them to be exceedingly fine specimens and can cordially recommend them to our friends. Mr. Eggert offers for sale or exchange a selection from a list of about 500 species collected in the vicinity of St. Louis. He allows those who wish to buy, the privilege of choosing from the list such plants as they desire to obtain, at the rate of 6 cents per species for full and complete specimens. Dr. George Engelmann is given as reference. The list may be procured by addressing H. Eggert, 918 Wash St., St. Louis, Mo.

In a note from Prof. M. W. Harrington, written from Leipzig, Germany, the following is of general interest. "The great German botanist, Prof. Hofmeister, died a few days ago. The two parts of his *Handbuch der Physiologischen Botanik*, promised for 1876, were not prepared and the book must therefore remain unfinished. To those who have possessed the earlier parts—some already 10 years old—this must be a source of great regret.

ALBINISM IN PLANTS.—In the Oct. No. of the BULLETIN I see a notice of a white *Lobelia syphilitica*, L. In confirmation of this fact, I will state that I have found two